

GlcPase was not inhibited under euglycemic hyperinsulinic conditions (group I), and was significantly inhibited under hyperglycemic conditions (G1 and G2). Our data strongly suggest that hyperglycemia, but not hyperinsulinemia, could account for the inhibition of liver Glc6Pase activity during the postprandial period. This phenomenon might play an important role in the suppression of HGP in this situation.

The expression of ob gene is not acutely regulated by insulin and fasting in human abdominal subcutaneous adipose tissue. H Vidal¹, D Aubœuf¹, P de Vos³, B Staels³, JP Riou^{1,2}, J Auwerx³, M Laville^{1,2} (¹ Inserm U 449, faculté de médecine René Laënnec, 69373 Lyon cedex 08; ² Centre de recherche en nutrition humaine de Lyon, Hôpital Edouard-Herriot, 69472 Lyon cedex; ³ Inserm U 325, Institut Pasteur, rue Professeur Calmette, 59019 Lille, France).

In rodents, the expression of leptin, the product of ob gene, is increased by insulin and decreased by fasting [Saladin et al (1995), *Nature* 377, 527-529; Trayhurn et al (1995), *FEBS Lett* 368, 488-490; Frederich et al (1995), *J Clin Invest* 96, 1658-1663]. In the present work, we investigated the regulation of ob gene expression in human abdominal subcutaneous adipose tissue using a reverse transcription-competitive PCR method to quantify the mRNA level of leptin. Leptin mRNA level was highly correlated ($r = 0.9$) with the body mass index of 26 subjects (12 lean, seven non insulin-dependent diabetic and seven obese patients). The effect of fasting on ob gene expression was investigated in ten subjects maintained on a hypocaloric diet (250 Kcal/day) for 5 days. While their metabolic parameters significantly changed (decrease in insulinemia, glycemia and resting metabolic rate and increase in plasma ketone bodies), the caloric restriction did not modify significantly the leptin mRNA

level in the adipose tissue. To verify whether insulin acutely regulates ob gene expression, six lean subjects underwent a 3 h euglycemic hyperinsulinemic ($846 \pm 138 \text{ pmol/L}$) clamp. Leptin and Glut 4 mRNA levels were quantified in adipose tissue biopsies taken before and at the end of the clamp. Insulin infusion produced a significant 3-fold increase in Glut 4 mRNA while leptin mRNA was not affected. It is concluded that ob gene expression is not acutely regulated by insulin or by metabolic factors related to fasting in human abdominal subcutaneous adipose tissue. Our data question therefore the role of leptin as a tightly controlled satiety factor in human but rather suggest that leptin signals the size of the adipose tissue deposit.

Glucose inhibits expression of the cytosolic aspartate aminotransferase gene in 3T3-F442A adipocytes: involvement of the promoter-regulatory region. E Pleegautier¹, M Aggerbeck², R Barouki², C Forest¹ (¹ Centre de recherche sur l'endocrinologie moléculaire et le développement, CNRS, 9, rue Jules-Hetzel, 92190 Meudon; ² Inserm U 99, Hôpital Henri-Mondor, 51, av Maréchal de Lattre-de-Tassigny, 94010 Créteil, France).

Aspartate aminotransferase (AspAT) is an ubiquitous enzyme involved in the malate-aspartate shuttle, urogenesis, gluconeogenesis and amino-acid metabolism. Two isoforms of the enzyme have been described: a cytosolic one (cAspAT) and a mitochondrial one (mAspAT). Although cAspAT is expressed at similar levels in all tissues, it is specifically regulated by glucocorticoids and glucagon in liver [Pavé-Preux et al (1988), *J Biol Chem* 263, 17459-17466] in which the enzyme is involved in gluconeogenesis. Regulation of cAspAT gene expression has been extensively studied in Fao hepatoma cells. In such cells, the rate of cAspAT gene transcription is increased by