

Involvement of voltage-dependent calcium channels (VDCC) in the action of GnRH on GtH release in common carp (*Cyprinus carpio* L): comparison with K⁺ action

T Mikolajczyk^{1, 2*}, C Weil¹, P Epler², B Breton¹

¹ INRA, Laboratoire de Physiologie des Poissons, Campus de Beaulieu, 35042 Rennes, France;
² Agricultural Academy, Department of Ichthyobiology & Fisheries, 30149 Crakow, Mydlniki, Poland

(Received 9 January 1990; accepted 20 July 1990)

Summary — The involvement of different types of voltage-dependent calcium channels (VDCC) in the stimulatory action of GnRH (in comparison with K⁺) on maturational gonadotropin (GtH) release was investigated using superfused carp pituitary cells. The action of these 2 stimulants was not modified either by D600 or nifedipine (drugs blocking L-type of VDCC). Cadmium (Cd²⁺), which blocks all types of VDCC indifferently, provoked a dose-dependent stimulation of GtH secretion. Cd²⁺ action was not altered by addition of sGnRH in any of the doses. Similar results were obtained using K⁺ as a secretagogue, but only the highest dose of Cd²⁺ (200 µmol/l) was able to completely block K⁺ action. Low doses (0.1 and 1 µmol/l) of the L-type VDCC activator BAY-K8644 did not change basal GtH secretion and had no effect on sGnRH-stimulated GtH secretion. Surprisingly, doses (10 µmol/l and higher) of BAY-K8644 evoked dose-dependent inhibition of GtH secretion. On the other hand, a higher concentration (20 µmol/l) of nifedipine provoked a stimulation of GtH release.

Our results indicate that the stimulatory action of GnRH and K⁺ involves activation of a certain type of cadmium-sensitive VDCC (probably T- or N-type VDCC) whereas dihydropyridine and di-phenylalkylamine sensitive VDCC (L-type VDCC) does not participate in this phenomenon. The inhibitory action of BAY-K8644 and, on the other hand, the stimulatory action of nifedipine indicate that L-type VDCC probably play a role in other physiological pathways regulating GtH release in carp.

GnRH / GtH / calcium channel / carp

Résumé — Implication de canaux calciques dont l'action dépend du voltage (VDCC) dans l'action du GnRH sur la libération de la GtH chez la carpe commune (*Cyprinus carpio* L): comparaison avec l'action de K⁺. Le rôle de différents types de canaux calciques dépendants du voltage (VDCC) dans l'action du GnRH sur la stimulation de la sécrétion de gonadotropine (GtH) par des cellules dispersées d'hypophyse de carpe a été étudié dans un système de perfusion. Le mécanisme d'action du GnRH a été comparé à celui du potassium (K⁺). L'action de ces 2 facteurs n'est modifiée ni par le D-600 ni par la nifedipine qui bloque les VDCC de type L. Le cadmium (Cd²⁺) qui bloque indifféremment tous les VDCC induit une stimulation dont l'intensité dépend de la sécrétion du GtH, qui n'est pas modifiée en présence de GnRH et ce quelle que soit la concentration de Cd²⁺. Des résultats comparables sont obtenus avec K⁺, mais seules de fortes concentrations de Cd²⁺ (200 µmol/l) bloquent l'action de K⁺. De faibles doses de BAY-K8644, activateur des canaux de type L, ne modifient ni la sécrétion basale de GtH, ni l'action du GnRH, alors que de fortes doses (10 µmol/l) induisent une inhibition dont l'intensité dépend de la sécrétion de GtH. Par ailleurs la nifedipine à 20 µmol/l stimule elle-même la sécrétion de GtH. Ces résultats montrent que l'action stimulante du

* Correspondence and reprints.

GnRH et de K⁺ implique l'activation de certains types de VDCC sensibles au cadmium, sans doute de type T ou N, les canaux sensibles aux dihydropyridine et diphényloalkylamine, de type L, n'étant pas mis en jeu dans ces phénomènes. Les actions inhibitrices du BAY-K8644, et stimulatrice de la nifedipine, montrent cependant que les canaux de type L jouent sans doute un rôle dans d'autres mécanismes régulant la sécrétion de GtH.

GnRH / GtH / canaux calciques / carpe

INTRODUCTION

It is well established that calcium ions (Ca²⁺) play the role of second messenger in GnRH action in mammals (Conn *et al*, 1987; Huckle and Conn, 1988). There are more and more reports indicating that GnRH action on GtH release is also a Ca²⁺-dependent process in fish (Jamaluddin *et al*, 1989; Levavi-Sivan and Yaron, 1989; Mikolajczyk *et al*, 1990). There is a general agreement that in higher vertebrates (birds and mammals), Ca²⁺ influx through voltage dependent calcium channels (VDCC) is responsible for prolonged LH secretion under GnRH stimulation. However, the spike phase of the secretory response to GnRH is independent of Ca²⁺ influx through VDCC (Hansen *et al*, 1987; Chang *et al*, 1986, 1988; Smith *et al*, 1987; Davidson *et al*, 1988; Smith *et al*, 1989). There could be an influx of Ca²⁺ through channels other than VDCC, the so-called receptor operated channels (ROC) (Conn *et al*, 1987; Davidson *et al*, 1988; Smith *et al*, 1989) or mobilization of Ca²⁺ from intracellular stores (Chang *et al*, 1986; Hansen *et al*, 1987). In fish literature there is also some controversy about the participation of VDCC in GnRH action. Jamaluddin *et al* (1989) observed in murrel a dose-dependent inhibition of GnRH-stimulated GtH secretion by D-600 (VDCC blocker) present in the culture medium. On the other hand, Van Asselt *et al* (1989) found that D-600 had no effect on spontaneous and GnRH-stimulated GtH release in African catfish, while such an inhibition

was observed when nifedipine (another class of VDCC blockers) was used.

We have shown an extracellular Ca²⁺ dependence of GnRH action on GtH release in carp (Mikolajczyk *et al*, 1990). The aim of the present study was to compare the action of sGnRH with K⁺ (elevated K⁺ evokes depolarization of cell membrane and activates VDCC) and to find out which type of VDCC participates in the stimulatory action of these 2 secretagogues on GtH release in carp.

MATERIALS AND METHODS

Animals

Experiments were conducted for 2 consecutive years (1988 and 1989) during the natural reproductive period of carp *ie* from April to July. Sexually mature female carp breeders weighing 5.5–11.5 kg obtained from the Heyman Fish Farm were kept in natural ponds and then transported to the laboratory. They were placed in a thermoregulated recirculating water system and acclimated at 18 °C for at least 10 d before being used in experiments. They were exposed to a controlled photoperiod (16L: 8D) and fed *ad libitum* with pelleted food (Aqualim).

Cell preparation and perfusion system

Cell preparation and the perfusion system were as previously described (Mikolajczyk *et al*, 1990). Briefly, pituitaries (1 pituitary for each perfusion) were dispersed using collagenase

(0.1%) (Boehringer Mannheim) in a medium (MEM-Eagle, Gibco) buffered with 15 mmol/l Hepes and 9 mmol/l sodium bicarbonate and supplemented with 1% BSA (RIA grade, Sigma). The mixture was incubated at 18 °C for 6 h, and dispersion was achieved mechanically. Cells were harvested by centrifugation, washed twice, mixed with Bio-Gel P-2 (Biorad) and placed in thermoregulated perfusion columns ($\pm 10^6$ cells per column). Five columns could be perfused at the same time. The cells were perfused first for 18 h at a flow rate of 4 ml/h with the above medium supplemented with 2% of a serum substitute, Ultrosor-SF (IBF), penicillin (100 U/ml) and streptomycin (10 µg/ml) (Gibco). The ultrosor and antibiotics were then withdrawn and the flow rate was gradually increased up to 15 ml/h. After 90 min of perfusion at a flow rate 15 ml/h, the first pulse of the drugs was administered. Fractions were collected every 7.5 min before drug administration and every 2.5 min during and 15 min after drug application. Details concerning each perfusion are given in the figure legends.

Drugs

Salmon GnRH (sGnRH) (Bachem), veratridine (Sigma) and cadmium chloride were dissolved directly in the perfusion medium prior to use. Methoxyverapamil (D-600), nifedipine (Sigma) and BAY-K8644 (RBI) were dissolved in ethanol and then in the perfusion medium. The amount of ethanol in the medium never exceeded 0.15%. The same amount of carrier was always present in the perfusion medium passing through control columns and had no measurable effect on GtH release. In experiments in which the effect of K⁺ depolarization was tested, the amount of NaCl in the test medium was reduced accordingly so that the total concentration of KCl and NaCl remained constant.

GtH determination and calculations

GtH levels were determined using a specific RIA developed by Breton *et al* (1971).

Profiles of GtH secretion are presented as a mean percentage of the basal GtH secretion lev-

el (bl). Basal level of GtH release (100%) was calculated as the mean of the 4 points (fractions) directly preceding first drug application. Fluctuations of GtH levels during the period directly preceding all drug administration never exceeded 20% of the basal GtH secretion level. Fluctuations of GtH levels greater than 20% were considered as significant. Differences in GtH secretion rate between the control and experimental columns during the stimulation period were calculated using Student's *t*-test for variation analysis.

RESULTS

Effect of organic VDCC blockers on GnRH- and K⁺-stimulated GtH release

As shown in figure 1A, two 15 min pulses of sGnRH (10^{-7} mol/l) evoked in the control columns a significant increase in GtH secretion (300% bl). In the experimental columns, the infusion of 2 µmol/l of VDCC blocker – D-600 (interfering mostly with L-type VDCC) had no effect on sGnRH-stimulated GtH release. A ten times higher (20 µM) concentration of D-600 had no effect on sGnRH action (fig 1B) either.

At 2 and 20 µmol/l concentrations, nifedipine, a specific L-type VDCC blocker had no effect on sGnRH-stimulated GtH release (448 and 310% bl) when administered in experimental columns in comparison with the control columns receiving sGnRH alone (432 and 350% bl) (fig 2). During the infusion of 20 µmol/l of nifedipine a significant increase ($P < 0.01$) in GtH secretion was observed (300% bl).

A 12.5-min application of KCl (60 mM of K⁺) resulted in a sharp increase (342 and 218% bl) in GtH secretion (fig 3 and data not shown). The amplitude and profile of the secretory response of the cells to elevated K⁺ was similar to that caused by sGnRH (figs 1, 2 and 5). No significant dif-




Fig 1A. Control columns (■) ($n = 5$) received two 15-min pulses of 10^{-7} M sGnRH (RH). Experimental columns (●) ($n = 5$) received the same pulses of sGnRH but the second one in the presence of $2 \mu\text{mol/l}$ of D-600. D-600 was present in the perfusion chamber 30 min before sGnRH application as indicated. Mean basal GtH level of all the columns was 31.5 ± 12.5 ng/ml. Data from 2 separate experiments. **B.** Eight columns (data from 2 experiments) received two 15 min pulses of 10^{-7} mol/l sGnRH (RH). The first one together with $20 \mu\text{mol/l}$ of D-600 as indicated. D-600 was present in the perfusion chamber 30 min before sGnRH application. Mean basal GtH level was 18.3 ± 8.3 ng/ml.

ferences were observed between the control columns receiving pulses of K^+ (60 mmol/l) and the experimental columns receiving K^+ in the presence of nifedipine ($2 \mu\text{mol/l}$) or D-600 ($2 \mu\text{mol/l}$) (fig 3).

Effect of Cd^{2+} , an inorganic VDCC blocker, on spontaneous, sGnRH- and K^+ -stimulated GtH release

A 30-min administration of graded doses of CdCl_2 (20 , 100 and $200 \mu\text{mol/l}$ of Cd^{2+}) resulted in a dose-dependent increase in GtH secretion (80 , 110 and 170% bl respectively) (fig 4). In the case of higher concentrations of Cd^{2+} (100 and $200 \mu\text{mol/l}$), the profile of GtH secretion exhibited a




Fig 2. Control columns (■) ($n = 7$) received two 15-min pulses of 10^{-7} M sGnRH (RH). Experimental columns (●) ($n = 8$) received the same pulses of sGnRH in the presence of 2 and $20 \mu\text{mol/l}$ of nifedipine as indicated. Nifedipine was applied 30 min before pulse of sGnRH. Mean basal GtH level of all the columns was 28.1 ± 10.1 ng/ml (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$) Similar results were obtained in 3 separate experiments.




Fig 3. Control columns (■) ($n = 7$) received two 12.5-min pulses of K^+ (60 mM). Experimental columns (●) ($n = 8$) received the same pulses of K^+ but in the presence of nifedipine or D-600 as indicated. Nifedipine and D-600 were present in the perfusion chamber 10 min before K^+ application. Mean basal level of all the columns from 3 separate experiments was 36 ± 20.7 ng/ml.

characteristic biphasic response: first phase during the presence of Cd^{2+} in the perfusion chamber and second phase after Cd^{2+} withdrawal (figs 4, 5A and 6). When sGnRH (10^{-7} mol/l) was infused during the application of Cd^{2+} (20 and $200 \mu\text{mol/l}$) no changes in GtH secretion were obtained in comparison with the control




Fig 4. Graded doses of Cd²⁺ were administered as 25-min pulses as indicated. Data represent a mean of 12 columns (3 experiments). Mean basal GtH level was 43.4 ± 17.8 ng/ml.

columns receiving Cd²⁺ alone (fig 5A). To check if Cd²⁺ had any side effect on sGnRH action (experiment presented in fig 5B) a similar pulse of sGnRH (alone) was applied 60 min after Cd²⁺ withdrawal after joint administration of Cd²⁺ (20 μ mol/l) and sGnRH (10^{-7} μ mol/l). This resulted, in both the experimental and the control columns, in the stimulation of GtH secretion, having a similar potency in a manner similar to earlier experiments.

The results presented in figure 6 show that 60 mM of K⁺ modified ($P < 0.05$) a secretory response of the cells to 100 μ mol/l concentration of Cd²⁺ in comparison with the control columns, while no differences were observed between the control and experimental columns during K⁺ stimulation when 200 μ mol/l of Cd²⁺ was applied.

Effect of organic activators of VDCC on GtH release

Veratridine (an organic membrane depolarizer) at a wide range of concentrations (0.1 to 10 μ mol/l) had no effect on GtH release (data not shown); a weak increase in GtH secretion (30–50% bl) was observed when much higher doses were used (50




Fig 5A. Five control columns (■) as well as 5 experimental columns (●) were exposed to 2 graded doses of Cd²⁺ (20 and 200 μ mol/l) for 30 min. In the experimental columns two 15-min pulses of sGnRH (RH) (10^{-7} mol/l) were administered 10 min after the beginning of Cd²⁺ application as indicated. Mean basal GtH level of all the columns from 2 separate experiments was 25.1 ± 15.0 ng/ml. **B.** Control columns (■) ($n = 5$) received a 30-min pulse of 20 μ mol/l of Cd²⁺ and 80 min later a 15 min pulse of sGnRH (10^{-7} mol/l) (RH). Experimental columns (●) ($n = 5$) received the same set of pulses as control columns but during the Cd²⁺ administration a 15-min pulse of sGnRH (10^{-7} mol/l) (RH) was applied 10 min after the beginning of Cd²⁺ infusion as indicated. Mean basal GtH level of all the columns was 42.3 ± 18.6 ng/ml. Data from 2 separate experiments.

and 100 μ mol/l respectively) (fig 7). The administration of BAY-K8644 (dihydropyridine, specific activator of L-type VDCC) in concentrations ranging from 0.01 to 1 μ mol/l caused no changes in GtH secretion (data not shown). BAY-K8644 at a concentration of 0.1 μ mol/l also had no effect on sGnRH-stimulated GtH secretion (data not shown). When BAY-K8644 was applied in concentrations of 10, 20 and 40 μ mol/l it




Fig 6. Control columns (■) ($n = 7$) were exposed to graded doses of Cd^{2+} in pulses lasting 30 min. In experimental columns (●) ($n = 8$) two 15-min pulses of K^+ (60 mM) were given 10 min after the beginning of Cd^{2+} application as indicated. Mean basal GtH level of all the columns from 3 separate experiments was 19.4 ± 6.1 ng/ml (See fig 2 for other details).

resulted in dose-dependent inhibition of GtH secretion – 28, 34 and 45% bl respectively (fig 8).

DISCUSSION

In recent years significant progress has been made in detailed characterization of the mechanism of GnRH action on gonadotropin release, mainly in rats and chickens; however there is more and more data concerning the mechanism of GnRH action in other groups of vertebrates, especially in fish (Levavi-Sivan and Yaron, 1989; Chang *et al*, 1989; Jamaluddin *et al*,




Fig 7. Eight columns (data from 2 experiments) received 2 pulses of graded doses of veratridine lasting 30 min as indicated. Mean basal GtH level was 26.3 ± 8.5 ng/ml.




Fig 8. Cells received three 15-min pulses of graded doses of BAY-K8644 as indicated. Data represent results from 4 separate experiments (16 columns). Mean basal GtH level was 29.8 ± 18.3 ng/ml.

1989; Van Asselt *et al*, 1989; Mikolajczyk *et al*, 1990). It was shown in mammals, birds, and recently in fish that an increase in intracellular Ca^{2+} concentration is an essential step in GnRH action and that Ca^{2+} plays the role of second messenger in GnRH action (Conn *et al*, 1987; Huckle and Conn, 1988). We also proposed such a role for extracellular Ca^{2+} in GnRH action on GtH release in common carp (Mikolajczyk *et al*, 1990). The aim of the present study was to determine which type of calcium channel is responsible for Ca^{2+} influx during GnRH stimulation. The results presented here indicate that the stimulatory action of K^+ on GtH release resembles GnRH action, indicating that the mechanism of GnRH action may involve the activation of voltage-dependent calcium channels (VDCC). It has been shown (Nowycky *et al*, 1985) that in neuronal tissue, 3 types (L, N and T) of VDCC exist. Each type of channel exhibits different electrophysiological and pharmacological properties (Miller, 1987). It is also possible that in pituitary cells several types of calcium channel exist, as reported by Armstrong and Matteson (1985) in a clonal line of rat pituitary cells. In our experiments we used drugs from 2 chemical classes as VDCC blockers: dihydropyridine (DHP) and diphenyl-

alkylamine (DPA). Nifedipine (DHP) specifically blocks the L-type VDCC (so-called DHP-sensitive). D-600 (DPA) also blocks the L-type VDCC. Since, in these experiments, nifedipine and D-600 did not inhibit either K⁺ or GnRH action, it is clear that these 2 secretagogues (K⁺ and GnRH) activate other types of VDCC than DHP- and DPA-sensitive (L-type) VDCC. To verify the role of other types of VDCC we applied cadmium ions (Cd²⁺) which block indifferently all types of VDCC. Surprisingly, all the doses of Cd²⁺ used evoked a stimulation of GtH release. In this situation, to see the effect of Cd²⁺ on sGnRH and K⁺ action we were obliged to look for a modification of Cd²⁺ stimulatory action by sGnRH and K⁺. Using sGnRH as a secretagogue it was impossible to modify the profile and amplitude of the secretory response of the cells to Cd²⁺ (figs 5A and 5B). However, elevated K⁺ significantly changed the secretory response to 20 µmol/l Cd²⁺ (data not shown) and to 100 µmol/l Cd²⁺ (fig 6). When Cd²⁺ was applied at a concentration of 200 µmol/l, no changes in GtH secretion were observed when K⁺ was introduced. When these results were combined, it was concluded that Cd²⁺ inhibits and blocks sGnRH and K⁺ action. It was however necessary to use a 10-times higher dose of Cd²⁺ to completely block the K⁺ action than was the case for sGnRH. This is not surprising since elevated K⁺ causes a total depolarization of the cell membrane and is probably a much stronger stimulatory factor than sGnRH. One of the possible explanations of the stimulatory effect of Cd²⁺ could be that different VDCC blocked by Cd²⁺ have a different influence on GtH release. For example, the blockade of a certain type of VDCC could inhibit GnRH and K⁺ action while at the same time, blockade of another type of VDCC could stimulate GtH release. This hypothesis is strongly supported by results obtained using specific drugs acting on L-type of VDCC. At low

doses nifedipine and BAY-K8644 had no effect on spontaneous and GnRH-stimulated GtH release. However at higher doses (10 µmol/l and more) nifedipine (channel blocker) stimulated GtH secretion with high potency and BAY-K8644 (channel activator) inhibited GtH secretion. Combined, these results suggest that during the action of Cd²⁺ there was a blockade of the L-type of VDCC and, as in the case of nifedipine, there was a stimulation of GtH release. At the same time there was also a blockade of N and T channels provoking the inhibition of K⁺ and GnRH action. In addition, a biphasic secretory response of the cells to Cd²⁺ also tends to support our hypothesis. Each time Cd²⁺ was applied, GtH stimulation was stronger at the end of the Cd²⁺ pulse than during Cd²⁺ administration. This suggests that the first phase of the response to Cd²⁺ was caused by a blockade of L-type VDCC and the second phase (after Cd²⁺ withdrawal) provoked by a rise in the blockade of, and/or the reactivation of N- and/or T-type VDCC. The fact that veratridine (an organic depolarizer) was able to stimulate GtH release only at very high doses and with very low potency indicates that this drug did not activate an exact type of VDCC responsible for GtH release. Another possible explanation of the surprising effect of NFD and BAY-K8644 could be their nonspecific action when used at a high concentration (> 10 µmol/l). Drouva *et al* (1988) observed an inhibition of PRL and GH secretion from rat pituitary cells evoked by 100 µmol/l BAY-K8644, whereas concentrations from 0.01 up to 10 µmol/l were stimulatory. On the other hand, other data from the same laboratory (Enjalbert *et al*, 1988) show that BAY-K8644 at a concentration of 100 µmol/l is less effective than at 10 µmol/l in stimulating PRL secretion but has no inhibitory effect on PRL release. Chang *et al* (1988) have also shown that BAY-K8644 at concentrations from 1 µmol/l to 10 µmol/l

significantly increased basal LH release and enhanced LH responses to GnRH.

Our results concerning the effect of D-600 and nifedipine differ however from results obtained by other investigators in fish. Jamaluddin *et al* (1989) using static pituitary cell culture found a dose-dependent inhibition of GnRH-stimulated GtH release in murrel using D-600. On the other hand Van Asselt *et al* (1989) using perfused catfish pituitary fragments found no effect of D-600 on GnRH-stimulated GtH secretion whereas GnRH action was inhibited by nifedipine. In our opinion such differences are provoked by different experimental approaches. It could be possible that in static cultures, only the prolonged phase of secretory response to GnRH was monitored as well as in the case of perfusion of pituitary fragments (Van Asselt *et al*, 1989) where fractions were rarely collected (10 min) to find any effect on the first phase of the secretory response. In any case, results obtained on pituitary fragments are not so convincing since it is impossible to determine the level of drug action and to avoid possible interactions inside whole pituitary tissue.

In mammals and birds it is well established that in the prolonged phase of LH secretion under GnRH stimulation there is an influx of extracellular Ca^{2+} via voltage-dependent calcium channels (VDCC). It was also shown that the L-type of VDCC (DHP- and DPA-sensitive) is responsible for this influx since specific L-type VDCC blockers inhibited or blocked the secretory response to GnRH (Hansen *et al*, 1987; Smith *et al*, 1987; Chang *et al*, 1988; Davidson *et al*, 1988; Smith *et al*, 1989). If the first (spike) phase of the secretory response to GnRH is concerned there is some controversy about the pathways as well as the sources of the Ca^{2+} . Chang *et al* (1986), Hansen *et al* (1987) proposed the possible participation of Ca^{2+} mobilized

from intracellular stores, whereas Conn *et al* (1987); Davidson *et al* (1988) and Smith *et al* (1989) found that there was an influx of extracellular Ca^{2+} via an other than VDCC, probably via so-called receptor operated channels (ROC). This second hypothesis is strongly supported by the findings of Mason and Waring (1985, 1986). They found that in bovine gonadotropes GnRH does not depolarize the cell membrane which is necessary to activate VDCC. On the contrary, Croxton *et al* (1988) have shown that in rat gonadotropes GnRH induces oscillatory membrane currents and this could be blocked by D-600. This finding tends to support the hypothesis of the intracellular origin of Ca^{2+} in GnRH action. These findings, together with the fact that several types of calcium channels exist could be an explanation for the striking differences among the results obtained on this subject.

CONCLUSION

The present findings indicate that GnRH action in carp is probably a voltage-dependent process involving activation of Cd^{2+} sensitive VDCC. These channels are probably T- or N-type of VDCC. DHP- and DPA-sensitive VDCC (L-type) does not participate in GnRH and K^+ action. Activation of L-type VDCC leads to inhibition and their blockade to stimulation of GtH release. This could be part of other physiological pathways regulating GtH secretion in carp. In fish, dopamine (DA) was shown to be a gonadotropin releasing inhibitory factor (GRIF), and application of DA antagonists (pimozide, domperidone) stimulates GtH secretion and strongly potentiates the action of GnRH (Peter *et al*, 1986). It has been shown that neuropeptide Y (NPY) also had a direct effect on GtH secretion in fish, and that its action depended on the

stage of sexual maturity of the fish (*ie* inhibitory action in vitellogenic animals, and stimulatory action in freshly ovulated fish) (Breton *et al.*, 1989, 1990). From our results it can be supposed that the inhibition of GtH release caused by activation of L-type VDCC, reflects the action of DA or other drugs inhibiting the secretion of GtH. It could also be speculated that Ca²⁺ serves as a second messenger for both DA and GnRH, but its different transport through the cell membrane (different types of calcium channels) and probably different intracellular Ca²⁺ receptors lead finally to a different effect on GtH release. This hypothesis is currently under investigation.

ACKNOWLEDGMENTS

The authors extend their thanks to M Gerber for the transport and care of fish. This study was supported by INRA, France.

REFERENCES

- Armstrong CM, Matteson DR (1985) Two distinct populations of calcium channels in a clonal line of pituitary cells. *Science* 227, 65-67
- Van Asselt LAC, Goos HJTh, Van Dijk W, Braas J (1989) Role of calcium ions in action of gonadotropin-releasing hormone on gonadotropin secretion in the African catfish, *Clarias gariepinus*. *Gen Comp Endocrinol* 76, 46-52
- Breton B, Kann G, Burzawa-Gerard E, Billard R (1971) Dosage radio-immunologique d'une hormone gonadotrope de carpe (*Cyprinus carpio* L.). *CR Acad Sci (Paris) Ser D* 272, 1515-1517
- Breton B, Mikolajczyk T, Danger JM, Gonnet F, Saint-Pierre S, Vaudry H (1989) Neuropeptide Y (NPY) modulates *in vitro* gonadotropin release from rainbow trout pituitary glands. *Fish Physiol Biochem* 7, 77-83
- Breton B, Mikolajczyk T, Weil C, Danger JM, Vaudry H (1990) Studies on the mode of action of neuropeptide Y (NPY) on maturational gonadotropin (GtH) secretion from perfused rainbow trout pituitary glands. *Fish Physiol Biochem* 8, 339-346
- Chang JP, McCoy EE, Graeter J, Tasaka K, Catt KJ (1986) Participation of voltage-dependent calcium channels in the action of gonadotropin-releasing hormone. *J Biol Chem* 261, 9105-9108
- Chang JP, Stojilkovic SS, Graeter JS, Catt KJ (1988) Gonadotropin-releasing hormone stimulates luteinizing hormone secretion by extracellular calcium-dependent and independent mechanism. *Endocrinology* 122, 87-97
- Chang JP, Freedman GL, De Leeuw R (1989) Participation of arachidonic acid metabolism in gonadotropin-releasing hormone stimulation of goldfish gonadotropin release. *Gen Comp Endocrinol* 76, 2-11
- Conn PM, McArdle CA, Andrews V, Huckle WR (1987) The molecular basis of gonadotropin-releasing hormone (GnRH) action in the pituitary gonadotrope. *Biol Reprod* 36, 17-35
- Conn PM, Staley DD, Yasumoto T, Huckle WR, Janovick JA (1987) Homologous desensitization with gonadotropin-releasing hormone (GnRH) also diminishes gonadotrope responsiveness to maitotoxin: a role for the GnRH receptor-regulated calcium ion channel in mediation of cellular desensitization. *Mol Endocrinol* 1, 154-159
- Croxton TL, Ben-Jonathan N, Armstrong WMCD (1988) Gonadotropin-releasing hormone induces oscillatory membrane currents in rat gonadotropes. *Endocrinology* 123, 1783-1791
- Davidson JS, Wakefield JK, King JA, Mulligan GP, Millar RP (1988) Dual pathways of calcium entry in spike and plateau phases of luteinizing hormone release from chicken pituitary cells: sequential activation of receptor-operated and voltage-sensitive calcium channels by gonadotropin-releasing hormone. *Mol Endocrinol* 2, 382-390
- Drouva SV, Rerat E, Bihoreau C, Laplante E, Rasolonjanahary R, Clauser H, Kordon C (1988) Dihydropyridine-sensitive calcium channel activity related to prolactin, growth hormone, and luteinizing hormone release

- from anterior pituitary cells in culture: interactions with somatostatin, dopamine, and estrogens. *Endocrinology* 123, 2762-2773
- Enjalbert A, Musset F, Chenard C, Priam M, Kordon C, Heisler S (1988) Dopamine inhibits prolactin secretion stimulated by the calcium channel agonist Bay-K-8644 through a pertussis toxin-sensitive G protein in anterior pituitary cells. *Endocrinology* 123, 406-412
- Hansen JR, McArdle CA, Conn PM (1987) Relative roles of calcium derived from intra- and extracellular sources in dynamic luteinizing hormone release from perfused pituitary cells. *Mol Endocrinol* 1, 808-815
- Huckle WR, Conn PM (1988) Molecular mechanism of gonadotropin-releasing hormone action. II. The effector system. *Endocr Rev* 9, 387-395
- Jamaluddin MD, Banerjee PP, Manna PR, Bhattacharya S (1989) Requirement of extracellular calcium in fish pituitary gonadotropin release by gonadotropin hormone-releasing hormone. *Gen Comp Endocrinol* 4, 190-198
- Levavi-Sivan B, Yaron Z (1989) Gonadotropin secretion from perfused tilapia pituitary in relation to gonadotropin-releasing hormone, extracellular calcium and activation of protein kinase C. *Gen Comp Endocrinol* 75, 187-194
- Mason WT, Waring DW (1985) Electrophysiological recordings from gonadotrophs. Evidence for Ca^{2+} channels mediated by gonadotropin-releasing hormone. *Neuroendocrinology* 41, 258-268
- Mason WT, Waring DW (1986) Patch clamp recordings of single ion channel activation by gonadotropin-releasing hormone in ovine pituitary gonadotrophs. *Neuroendocrinology* 43, 205-219
- Mikolajczyk T, Weil C, Epler P, Breton B (1990) Calcium ions as a mediator in GnRH action on gonadotropin release in the common carp (*Cyprinus carpio* L). *Reprod Nutr Dev* 30, 483-492
- Miller RJ (1987) Multiple calcium channels and neuronal function. *Science* 235, 46-52
- Nowycky MC, Fox AP, Tsien RW (1985) Three types of neuronal calcium channel with different calcium agonist sensitivity. *Nature* 316, 440-443
- Peter RE, Chang JP, Nahorniak CS, Omeljaniuk RJ, Sokolowska M, Shih SH, Billard R (1986) Interactions of catecholamines and GnRH in regulation of gonadotropin secretion in teleost fish. *Recent Prog Horm Res* 42, 513-548
- Smith CE, Wakefield I, King JA, Naor Z, Millar RP, Davidson JS (1987) The initial phase of GnRH-stimulated LH release from pituitary cells is independent of calcium entry through voltage-gated channels. *FEBS Lett* 225, 247-250
- Smith CE, Davidson JS, Millar RP (1989) Ba^{2+} stimulation of luteinizing-hormone release demonstrates two mechanisms of Ca^{2+} entry in gonadotrope cells. *Biochem J* 259, 217-221