

Microstructure of skeletal muscles of growing calves fed silage-based vs hay-based diets.

II. Fibre type distribution

B Kłosowski ^{1*}, K Bidwell-Porębska ²,
D Kłosowska ^{1*}, J Piotrowski ^{2**}

¹ Meat and Fat Research Institute, Warsaw, Laboratory of Meat Science, Bydgoszcz;

² Polish Academy of Sciences, Institute of Genetics and Animal Breeding,
Jastrzebiec, 05-551 Mroków, Poland

(Received 9 September 1991; accepted 22 April 1992)

Summary — As described in part I, samples of musculus longissimus dorsi, semimembranosus and semitendinosus were obtained post-slaughter from 2-week, 3-month and 10-month-old bull calves. The 2-week-old calves were fed milk only. All the remaining animals were fed grass silage or hay *ad lib* and a restricted amount of concentrate from 2 weeks of age onwards. Muscle fibres were differentiated according to Ziegen (1979) into fast-twitch glycolytic, fast-twitch oxidoglycolytic and slow-twitch oxidative fibres (FTG, FTO and STO, respectively). The percent distribution of individual types of fibres was estimated as related to the calves age and diet. The most numerous were always fast-twitch glycolytic fibres, the lowest values being observed in the 2-week-old calves. The effect of the diet on fibre percentage distribution in 3-month-old calves differed from that found in 10-month-old animals. This research suggests that a hay-free diet based on grass silage alters the microstructure of skeletal muscles, which thus might also affect the quality of meat.

calf / skeletal muscle fibre / silage / hay

Résumé — Microstructure des muscles squelettiques chez des veaux recevant de l'ensilage ou du foin. II. Pourcentage des fibres musculaires. Des échantillons de muscles longissimus dorsi, semimembranosus et semitendinosus furent collectés, comme décrit dans la partie I à la suite de l'abattage de veaux de 2 semaines, et de taurillons de 3 mois et de 10 mois. Les veaux âgés de 2 semaines avaient été nourris au lait, tandis que les veaux restants reçurent à volonté de l'ensilage de graminées ou du foin, ainsi qu'une dose limitée d'aliment concentré. Conformément à la méthode de Ziegen (1979), ont été distinguées les fibres musculaires glycolytiques et oxydatives glycolytiques se contractant rapidement, et les fibres oxydatives à contraction lente. Les plus nombreuses étaient toujours les fibres glycolytiques à contraction rapide – les valeurs les plus basses concernant les veaux de 3 mois différait de celle observée sur des taurillons de 10 mois. Ce travail suggère qu'une ration sans foin, à base d'ensilage de graminées, influence la microstructure des muscles squelettiques et pourrait donc agir sur la qualité de la viande.

veau / fibre de muscle squelettique / ensilage / foin

* Present address: Technical and Agricultural University, Faculty of Animal Science, Mazowiecka 28, 85-084 Bydgoszcz, Poland

** Correspondence and reprints

INTRODUCTION

The fibre is the basic unit of skeletal muscle and comprises 75 to 90% of the total muscle mass (Hegarty, 1971). It has been proved in domestic animals that the total muscle fibre numbers remain unchanged from birth or shortly before, and throughout the animal's life (Staun, 1963; Ashmore and Addis, 1972). Further growth of the muscle is a result of changes in the proportions between fibre types and of an increase in myofibre size (Ashmore *et al*, 1972; Kłosowski and Kłosowska, 1987).

It is well established in cattle that the proportions between muscle fibre types change with age and body weight (Cornforth *et al*, 1973; Johnston *et al*, 1975; Kłosowski and Kłosowska, 1988). However, few studies have been made on young, growing animals to determine the influence of different nutrition programmes. Meanwhile, despite the extensive use of silages in cattle nutrition, the effect of all-silage forage programmes on the organism is often controversial. The negative influence of organic acid concentrations and low pH on the digestive tract is generally accepted (Ballarini, 1974; Pellegrini *et al*, 1982a). Vitamin A, mineral and iodine deficiencies, known to be connected with silage nutrition (Hemken and Vandersall, 1967; Pellegrini *et al*, 1982a, b), may also cause "dystrophic regressive phenomena in the skeletal muscle fibres" (Pellegrini *et al*, 1982c) and probably influence the market and technological value of the meat. In part I of this study we have investigated the effect of silage-based or hay-based diets on the growth of fibres of selected skeletal muscles in calves (Kłosowski *et al*, 1992). Part II examines the percentage distribution of fibre types.

MATERIAL AND METHODS

The collecting, grouping, feeding and slaughtering of calves as well as calf performance have been described in part I of this study (Kłosowski *et al*, 1992). Following the paper cited, the calves were referred to as hay diet (HD, group I and IA) or silage diet (SD, group II and IIA)-fed animals, and as "younger" (3 months old: group I and II) or "older" (10–11 months old: group IA and IIA). A group of 5 calves (2 weeks old), referred to as "newborn", was also included.

Single tissue samples were taken approximately 1 h post-slaughter from musculus longissimus dorsi (longissimus lumborum – LD), musculus semimembranosus (SM) and musculus semitendinosus (ST). In part I the sampling, storing, sectioning, preparation, and staining procedures have been described (Kłosowski *et al*, 1992). Fast twitch glycolytic (FTG), slow twitch oxidative (STO) and fast twitch oxidoglycolytic (FTO) fibres were identified according to Ziegler (1979). The percent distribution of fibre types within each muscle was estimated in 10 primary bundles of fibres of each type (approximately 400 individual fibres per sample).

The significance of the relationship between calf age or diet and fibre distribution was calculated within the groups and for each muscle separately using the χ^2 test, suitable for calculations on percent values. The information thus obtained made it possible to evaluate the differences between the general structure of individual muscles within the age and nutrition groups.

RESULTS

The data referring to the daily nutrient intake and growth rate for all the age and feeding groups have been presented in part I of this study (Kłosowski *et al*, 1992).

Figure 1 indicates that the most numerous were the FTG fibres, constituting roughly 40–54% of all fibres, the lowest values being observed for newborn calves. In the case of LD muscle, the percentage of FTG fibres remained unchanged until the animals were 90 days old and in-




Fig 1. Percent distribution of fibre types (FTG: fast-twitch glycolytic, STO: slow-twitch oxidative, FTO: fast-twitch oxydoglycolytic) in individual muscles (LD: longissimus dorsi, SM: semimembranosus, ST: semitendinosus) as related to calf age. See table I for statistical estimations.

Table I. Significance of relationship (χ^2) between the calves age and percent distribution of fibre types presented in figure 1.

<i>Animals compared</i>	<i>Musculus longissimus dorsi</i>	<i>Musculus semimembranosus</i>	<i>Musculus semitendinosus</i>
Newborn and all younger calves	xx	xx	xx
Newborn and all older calves	xx	x	xx
All younger and all older calves	x	NS	xx

x: $P < 0.05$; xx: $P < 0.01$; NS: non significant.

creased slightly in the older calves. In the SM and ST muscles the percentage of FTG fibres increased already in the young-

er calves and remained fairly constant in the older animals. The percentage distribution of FTO fibres presented a picture con-




Fig 2. Percent distribution of fibre types (1: fast-twitch glycolytic, 2: slow-twitch oxidative, 3: fast-twitch oxidoglycolytic) in individual muscles (as in figure 1) of 3-month-old calves, as related to diet. 0.01 and 0.05: significance of relationship (χ^2); ns: non significant.

trary to that of the FTG: it was markedly highest in newborn, and lowest in younger calves, the differences related to age being most marked in the SM and ST muscles. The differences in percent distribution of STO fibres between age groups were the smallest and the least regular.

Using the χ^2 test for comparing newborn to all younger as well as all older calves (table 1), a highly significant relationship was demonstrated between calf age and the percent distribution of fibre types in each of the three muscles. How-

ever, when comparing the younger and older calves the relationship was less convincing, as it was found significant only for LD and ST muscles.

No significant effect of diet was proved when comparing feeding systems within younger calves (fig 2). However, in older calves (fig 3) such an effect was found to be significant for the SM and ST muscles. This was caused principally by the demonstrated differences in the percentage share of FTG and STO fibres between the nutrition groups.




Fig 3. Percent distribution of fibre types (as in figure 2) in individual muscles (as in figure 1) of 10–11-month old calves as related to diet. Statistical estimation as in figure 2.

DISCUSSION

The percentage distribution of individual fibre types corresponds with that reported by other authors (Melton *et al*, 1971; Hunt and Hendrick, 1977; Kłosowski and Kłosowska, 1984). The influence of age is also consistent with earlier investigations. In this study changes in the percentage distribution of fibres, in particular an increase of the share of fibres characterised by a glycolytic metabolism, were also related to age and live weight, and differed for different muscles (fig 1). This can be related to their various functions and non-identical growth patterns (Ashmore *et al*, 1972; Cornforth *et al*, 1973; Richmond and

Berg 1982). The differences in the percentage distribution, arising principally from an increase in the number of FTG fibres and a simultaneous decrease in the number of FTO fibres, could be attributed to a transformation between fibre types, as suggested by Ashmore *et al* (1972). The different developmental rate of LD as compared with SM and ST muscles (fig 1) was observed earlier by Kłosowski and Kłosowska (1988), who also did not observe any changes in the share of FTG fibres in the LD between 2 week- and 3 month-old calves.

The different effect of the diet on fibre distribution in younger vs older calves is interesting. Diet did not influence fibre type

distribution in younger calves (fig 2). However, in the older animals (fig 3), those fed hay had a higher percentage of FTG and FTO, and simultaneously a lower proportion of STO fibres; only in the case of the LD muscle were no differences observed between the 2 feeding groups. Moody *et al* (1980) when comparing the percentage distribution of fibres in the LD muscle also found no significant differences between animals receiving different diets. However, in their experiment the difference in the feeding regime lay in the presence or absence of pasture. Moreover, they only examined the LD muscle, paying no attention to the different growth rates which characterise different muscles.

In the present study, it is difficult to interpret the higher percentage of FTG and FTO fibres, and the lower percentage of STO fibres in the SM and ST muscles of the older HD fed animals. The older calves from both feeding groups were slaughtered at a constant live weight of approximately 250 kg (338 vs 305 days of age, for HD and SD group, respectively). However, the mean DM intake (3075 vs 4599 g/day) and GE (59.5 vs 86.9 MJ/day), and thus also the mean live weight gains (617 vs 713 g/day) were significantly lower in the HD as compared with the SD-fed calves. Thus the differences observed in the present distribution of fibres in the ST and SM muscles could be attributed to the different daily DM and GE intake in the 2 feeding groups, and consequently to different daily LW gains. The results reported by Johnston *et al* (1981) confirm the influence of energy concentration in the diet on muscle fibre distribution, but in reverse, the higher the energy level, the higher the percentage of fibres connected with the glycolytic metabolism. Such a discrepancy may be caused by the silage diet as such. In part I of the present investigation (Kłosowski *et al*, 1992), differences have also been observed in the fibre diameters be-

tween the 2 feeding groups, which could not be adequately explained by differences in nutrient intake or growth rate. This problem needs to be examined more thoroughly, all the more so as certain investigations have shown all silage diets to cause considerable histopathological changes, not only in skeletal muscles (Pellegrini *et al*, 1982c) but also in numerous internal organs of cattle (Pellegrini *et al*, 1982a, b; Slesareva and Soloveva, 1968). The present investigation suggests that such diets can also affect the percent distribution of fibres of skeletal muscles. Whether this may influence the final quality of meat remains an interesting question, especially as Rumsey *et al* (1987) reported silage feeding to affect various beef characteristics, while Ashmore *et al* (1972) demonstrated that meat quality can be related to changes in proportions between muscle fibre types.

REFERENCES

- Ashmore CR, Addis PB (1972) Prenatal development of muscle fiber types in domestic animals. *Proc Recip Meat Conf* 25, 211
- Ashmore CR, Tompkins G, Doerr L (1972) Postnatal development of muscle fiber types in domestic animals. *J Anim Sci* 34, 37-41
- Ballarini G (1974) La patologia alimentare da insilato di mais nei ruminanti. *Riv Zoot Vet* fasc 4, 355-370
- Cornforth DP, Schwartz WC, Cramer DA (1973) Growth and differentiation of muscle fibers in cattle. *J Anim Sci* 36, 1196
- Hegarty PVG (1971) Muscle fibre growth and development. *Proc Recip Meat Conf* 24, 319
- Hemken RW, Vandersall JH (1967) Feasibility of an all silage forage program. *J Dairy Sci* 50, 417-422
- Hunt MC, Hedrick HB (1977) Histochemical and histological characteristics of bovine muscles from four quality groups. *J Food Sci* 42, 578-582
- Johnston DM, Stewart DF, Moody WG, Boling J, Kemp JD (1975) Effect of breed and time of

- feed on the size and distribution of beef muscle fiber types. *J Anim Sci* 40, 613-620
- Johnston DM, Moody WG, Boling JA, Bradley NW (1981) Influence of breed type, sex, feeding systems and muscle bundle size on bovine fibre type characteristics. *J Food Sci* 46, 1760
- Kłosowski B, Kłosowska D (1984) Wpływ intensywności opasu na aktywność enzymatyczną i wielkość włókien mięsniowych buhajów rasy ncb. *Zesz Probl Post Nauk Roln* 267, 397-403 (in Polish; English summary)
- Kłosowski B, Kłosowska D (1987) Histochemical and histological characteristics of the bulls muscles during growth. Proc VIIIth Congr Polish Soc Vet Sci part III, 57-58 (in Polish; English summary)
- Kłosowski B, Kłosowska D (1988) Age related changes of the fiber characteristics in longissimus dorsi muscle of Lowland Black-and-White cattle. Vortrage gehaltete auf dem 4. Morphologie-Kolloquium des Wissenschaftsbereiches Anatomie, Histologie und Embriologie "Wachstum und Postnatale Entwicklung". Karl-Marx Univ, Leipzig, 23 June 1988, 109-113 (in German; English summary)
- Kłosowski B, Bidwell-Porębska K, Kłosowska D, Piotrowski J (1992) Microstructure of skeletal muscles of growing calves fed silage-based vs hay-based diets. I. Fibre diameters. *Reprod Nutr Dev* 32, 1-10
- Melton CC, Dikeman ME, Tuma HJ (1971) Histochemical studies of bovine longissimus muscle biopsies as related to meat quality and cuttability. *J Anim Sci* 33, 221
- Moody WG, Kemp JD, Mahyuddin M, Johnston DM, Ely DG (1980) Effect of feeding systems, slaughter weight and sex on histological properties of lamb carcasses. *J Anim Sci* 50, 249-255
- Pellegrini N, Renzoni G, Taccini E, Filippi Balestra G, Bagatti E (1982a) Effects of a silo maize on cows. Note 1. Anatomohistopathological modifications of the digestive system. *Ann Fac Med Vet Pisa* 35, 31-40 (in Italian; English summary)
- Pellegrini N, Renzoni G, Taccini E, Filippi Balestra G, Bagatti E (1982b) Effects of a silo-maize diets of cows. Note III. Histomorphological modifications of the thyroid, pancreas and adrenal glands. *Ann Fac Med Vet Pisa* 35, 49-58 (in Italian; English summary)
- Pellegrini N, Renzoni G, Taccini E, Filippi Balestra G, Bagatti E (1982c) Experimental evaluation of the visceral and organic dystrophic effects in calves fed with silo-maize only. Note V. Anatomohistopathological findings in the heart and skeletal muscle. *Ann Fac Med Vet Pisa* 35, 67-72 (in Italian; English summary)
- Richmond RJ, Berg RT (1982) Relative growth patterns of individual muscles in the pig. *Can J Anim Sci* 62, 575-586
- Rumsey TS, Bond J, Berry BW, Hammond AC, Dinius DA (1987) Performance and carcass characteristics of feedlot steers fed all silage diets. *Nutr Rep Int* 35, 847-862
- Slesareva EN, Soloveva VN (1968) Changes in the bone and udder of milking cows fed on silage throughout the year. *Izv Timir Selskokhoz Akad* 2, 172-183
- Staun H (1963) Various factors affecting number and size of muscle fibers in the pig. *Acta Agric Scand* 13, 293-322
- Ziegen J (1979) Combinations of enzyme-histochemical methods for differentiating of fibres types and evaluating the skeletal musculature. *Acta Histochem Jena* 65, 1, 34-40 (in German; English summary)