

released, leading to a prolonged and larger oxidation of exogenous glucose.

COLONIC MICROFLORA AND METABOLIC FUNCTIONS

Molecular mechanisms of butyrate action on HT-29 intestinal epithelial cell proliferation. S. Siavoshian, J.P. Segain, C. Cherbut, J.P. Galmiche, H.M. Blottière (Human Nutrition Research Center, CRI Inserm 95-08, CHU Hôtel-Dieu, Inra, 44035 Nantes cedex 01, France).

Sodium butyrate, a product of colonic bacterial fermentation, is able to inhibit cell proliferation and to induce the differentiation of colonic epithelial cells in culture. In a variety of cell systems, butyrate has been found to block cells in the G1 phase. D type cyclins appear early in the G1 phase and complex with cdk4 or cdk6. Then, they can bind to the unphosphorylated form of pRb, thus allowing E2F transcription factor to activate gene transcription. The p21 protein inhibits the action of G1/S cdk-cyclin complexes. The aim of our study was to investigate the mechanisms by which butyrate inhibits cell cycle progression toward S phase.

HT-29 cells were cultured in the presence or absence of increasing concentration of sodium butyrate (from 2 to 8 mM) for 24 h. Proteins were extracted, and cyclin D1, D3, E2F-1 and p21 expression were studied by western blotting. mRNA were extracted, and cyclin D1, D2 and p21 expression were studied by RT-PCR.

Butyrate inhibited cyclin D1 mRNA expression, without affecting its protein level. In contrast, butyrate stimulated cyclin D3 protein expression. We failed to detect any mRNA for cyclin D2 in HT-29 cells. Moreover, a dose-dependent decrease in E2F-1 expression was observed in HT-29 exposed to butyrate. In addition, after 6 h of incubation with

butyrate, p21 mRNA was detected and mRNA expression reached a plateau in between 12 and 24 h. At the protein level, no p21 was detected at 6 h. At 12 h, p21 was detected and the optimal detection was observed at 24 h.

Our results suggest that the inhibition of cell cycle progression by sodium butyrate may be explained by a modulation of cell cycle regulatory proteins such as cyclin D3 and p21.

Measurement of short chain fatty acids produced from resistant starch fermentation in the portal blood: A study in a pig model. L. Martin^a, H. Dumon^a, G. Lecannu^b, M. Champ^{b,c} (^aEcole nationale vétérinaire, Unité de nutrition et alimentation, BP 40706, 44307 Nantes cedex 03; ^bInra, Laboratoire de technologie appliquée à la nutrition, BP 71627, 44316 Nantes cedex 03; ^cCRNH groupe métabolisme, Hôtel Dieu, Place A. Ricordeau, 44093 Nantes cedex 01, France).

Colonic fermentation of resistant starch (RS) produces short chain fatty acids (SCFA) and an especially high proportion of butyric acid (*n*C4). This SCFA plays a specific role in the wholeness of colonic mucosa. In this study, we tested the hypothesis that the rate of starch fermentation could influence the production of *n*C4 and its utilization by the colonocytes in vivo.

Material and methods: Three experimental diets were formulated with three different resistant starches selected on the basis of their in vitro fermentation kinetic: raw potato starch (PoS), high amylose corn starch (HS) and retrograded high amylose corn starch (RHS). The animals ate 15 g of RS per experimental meal. Four pigs were fitted with two indwelling catheters (portal vein and carotid artery) and a flow probe (Transonic™) was placed around the portal vein. Results were analysed by ANOVA.