

Effects of paternal heat stress on the *in vivo* development of preimplantation embryos in the mouse

Bi-ke ZHU^{a*}, Brain P. SETCHELL^b

^a Faculty of Veterinary Science, University of Sydney, Camden, NSW 2570, Australia
^b Department of Animal Science, University of Adelaide, Waite Campus, Glen Osmond, SA 5064, Australia

(Received 2 February 2004; accepted 18 September 2004)

Abstract – The objective of this study was to examine the effect of paternal heat stress on the *in vivo* development of preimplantation embryos in the mouse. Synchronised B6CBF1 female mice were mated either to a control male mouse or to one that had been exposed at 7, 21 or 35 days previously, for 24 h to an ambient temperature of 36 ± 0.3 °C and $66 \pm 5.6\%$ relative humidity. Embryos were collected from the oviducts of mice at 14–16 h, 34–39 h or 61–65 h after mating or from the uterus at 85–90 h after mating and their developmental status was evaluated morphologically. The number of cells within blastocysts was also determined using bisbenzimidazole-propidium iodide staining. Paternal heat stress 7 days before mating reduced the proportion of embryos developing from 4-cell (4-C) to morulae (M), hatched blastocysts, total blastocysts and the number of inner cell mass (ICM) and trophectoderm (TE) cells in the blastocyst. Paternal heat stress 21 days prior to mating reduced the proportion of 2-C and 4-C to M embryos with no embryos developing to blastocysts. There were also increases in the number of 1-C and abnormal embryos recorded at this time. Paternal heat stress 35 days before mating decreased the proportion of 2-C embryos, expanded blastocysts and ICM and TE cells in the blastocyst. These results support previous work demonstrating that both the sperm in the epididymis and germ cells in the testis are susceptible to damage by environmental heat stress, with spermatocytes being the most vulnerable. This study also demonstrates that subtle effects on the male such as a short exposure to elevated environmental temperatures can translate to quite profound paternal impacts on early embryo development.

paternal heat stress / *in vivo* development / mouse preimplantation embryos

1. INTRODUCTION

In most mammals, testicular temperature is maintained between 2 and 7 °C lower than the core body temperature, and slightly higher than the scrotal temperature [1–4] for normal spermatogenesis. Any factors causing an increase in testicular temperature (e.g. high ambient temperature, local heat stress)

can damage spermatogenesis, which may result in decreased sperm numbers and reduced sperm motility in the ram [5] and boar [6], normal fertilisation, but an increase in embryonic death in the pig [6], mouse [7–9], rabbit [10, 11], sheep [12–17] and rat [18].

The quality of semen in man has also been reported to vary with the season and

* Corresponding author: bzhu7534@usyd.edu.au

occupation. Sperm concentration, total sperm count per ejaculate and motile sperm concentration have been reported to be lower in the summer than in the winter [19–22] and even mild heat can temporarily reduce the number of sperm ejaculated after two to three weeks [23]. The effect of heat stress on spermatogenesis may further extend time to pregnancy and decrease birth rate in women [24].

Early studies have shown that paternal heat stress reduces fertilisation rates [8] and the *in vivo* development of preimplantation embryos in the mouse [7]. The extent of the effect could be related to the time after heat-stress of the males [8, 18]. Setchell et al. [18] reported that local heating of the testis between 7 days and 35 days prior to mating reduces the size and weight of litters in the mouse. Our own studies have also indicated that the *in vitro* developmental rates of heat-sired preimplantation embryos begin to decrease in the embryos sired on 3 days after heat-stress, were the lowest when embryos were sired on 21 days after heat-stress, and then gradually returned to control levels when sired on 42 days [9]. Therefore, in the present study, we investigated the effect of paternal heat stress on the *in vivo* development of preimplantation embryos conceived at 7, 21 and 35 days after heating of the male to determine the stage at which the preimplantation embryos might be affected by paternal heat stress.

2. MATERIALS AND METHODS

Procedures used for all experimental animals were approved by the Animal Ethics Committee, at the University of Adelaide.

2.1. Heat stress of the male

Twelve fertile B6CBF1 male mice (3 to 4 months old) were placed in a psychrometric chamber set at 36 ± 0.3 °C and $65.8 \pm 5.6\%$ relative humidity and heated for 24 h, while another twelve comparable fertile F1 male mice were similarly kept at 21 °C as

a control group. This experiment was undertaken in triplicate.

2.2. Determination of sperm characteristics

The control and heat-stressed males were killed by cervical dislocation, 7, 21 and 35 days after the treatment. The epididymides were collected and the cauda epididymis was isolated and placed in a Falcon tube containing 2 mL of HTF medium (pH 7.4). The tissue of the cauda epididymis was minced by using scissors to release spermatozoa. The spermatozoa were allowed to swim up and incubated for 1 h in an atmosphere of 5% CO₂, 5% O₂ and 90% N₂ at 37 °C, the concentration and motility of spermatozoa were determined, using Microcell chambers (Conception Technologies, USA). For the determination of sperm concentration, 6 µL of the diluted semen were loaded into a Microcell chamber. An Olympus phase-contrast microscope fitted with an eyepiece reticle (Olympus Australia Pty. Ltd. Australia) was used to score sperm from 6 or 12 fields in each side of the chamber. Two hundred to three hundred spermatozoa were scored in each chamber. For the determination of motile sperm, semen was loaded into a prewarmed (37 °C) microcell chamber and placed on the heated (37 °C) microscope stage. The non motile and total motile (any type of movement) sperm were counted by looking through the reticle grid in the microscope eyepiece. Two hundred spermatozoa were scored in each Microcell chamber. The viability and morphology of spermatozoa were evaluated with a staining method of eosin-nigrosin [25]. Ten microliters of diluted semen stained with a solution containing 0.67% eosin (Sigma, Australia) and 5% nigrosin (Sigma, Australia) was applied to a slide which was kept on a warm plate at 37 °C and smeared using a coverslip (18 × 18 mm). The slides were left to dry on the warm plate for about 5 min. The percentage live (unstained, white in colour)/dead (stained, pinkish in colour) spermatozoa was determined by counting 200 sperm

cells in each of three prepared slides using an Olympus bright field microscope fitted with eyepiece reticles (Australia) (400×). The prepared slides were also used for morphological evaluation. Normal sperm showed no sign of defect but had the ideal sperm cell structure. Abnormal sperm showed defects with the head, midpiece or tail.

2.3. Oestrous synchronisation of female mice

Normal B6CBF1 female mice (2 to 3 months old) were placed in a room with no males present. After 28 days, a small cage containing male mice was placed on top of a large cage containing female mice so that the "male effect" would shorten and synchronise the female oestrous cycles [26, 27]. Approximately 30% of female mice were in oestrus on the third day following exposure to the male, and showed vaginal gaping, with vaginal tissues being lighter pink and less moist, and the striations being more pronounced.

2.4. Mating and collection of preimplantation mouse embryos

Oestrous female mice were mated with control or heat-stressed males. Each control male and heat-stressed male was caged with 2 oestrous females on 7, 21 and 35 days after heating of the male. The existence of a copulation plug the following morning, was taken as an indication of a successful mating.

Successfully mated mice were euthanised by cervical dislocation at specific intervals after mating. The mating time was defined as midnight (0.00 h) and female mice were allocated randomly for the collection of preimplantation embryos at the zygote, 2-cell, 4-cell to morula and blastocyst stages from the oviduct or the uterus at 14–16 h, 34–39 h, 61–65 h and 85–90 h after mating, respectively. The number of corpora lutea, the number of preimplantation embryos recovered and their stages of development were recorded for each female.

2.5. Morphological evaluation of preimplantation embryos

The quality of the individual embryos was determined by a method described previously [28], scoring normal, abnormal or unfertilised. The definitions applied were:

2.5.1. Normal

The embryo was spherical and symmetrical with cells of uniform size, colour and texture; the embryo was at an appropriate stage of development, i.e. the embryos recovered at 14–16 h, 34–39 h, 61–65 h and 85–90 h after mating were zygotes judged by the presence of the second polar body or pronuclei, 2-cell embryos, 4-cell embryos to morulae and morulae to blastocysts, respectively.

2.5.2. Abnormal

The embryos showed extruded blastomeres, a few cells of variable size, and contained numerous large vesicles.

2.5.3. Unfertilised

An ovulated ovum which did not contain the second polar body or pronuclei.

2.6. Differential fluorescent labelling of trophoblast (TE) and inner cell mass (ICM) in the blastocyst

The number of TE and ICM cells was determined by a method described previously [28]. Blastocysts were transferred from a HEPES-HTF medium [29] to acid Tyrode solution [30], under constant observation for 30–60 s, until the zona pellucida were completely dissolved. They were then washed 3 times in the HEPES-HTF medium and transferred into a CZB medium [31] containing 10% rabbit anti-mouse whole serum (M5774, Sigma, Australia), and then incubated at 37 °C for 15 min. The embryos were washed 3 times in the HEPES-HTF medium and transferred to the CZB medium containing 10% guinea pig serum complement (51639, Sigma, Australia) and incubated at 37 °C for 5–6 min. They were then

Table I. Semen characteristics of male mice, 7, 21 and 35 days after the heated treatment.

Groups*	C (N = 18)	H7 (N = 6)	H21 (N = 6)	H35 (N = 6)
Sperm concentration ($\times 10^6 \cdot \text{mL}^{-1}$)	13.5 \pm 0.37 ^a	9.1 \pm 0.58 ^b	6.5 \pm 0.76 ^c	10.2 \pm 0.37 ^b
Sperm motility (%)	68.5 \pm 2.13 ^a	44.2 \pm 5.23 ^b	30.5 \pm 2.89 ^c	54.2 \pm 3.74 ^b
Sperm viability (%)	69.6 \pm 1.12 ^a	62.5 \pm 2.05 ^b	36.7 \pm 1.63 ^c	66.7 \pm 1.26 ^a
Sperm abnormality (%)	6.5 \pm 0.46 ^a	18.1 \pm 1.39 ^b	38.3 \pm 1.06 ^c	12.5 \pm 0.76 ^d

* At each time point, 6 controlled and 6 heated males were killed for the determination of semen characteristics. The data obtained from 18 controlled males (C) were pooled together because there were no significant differences in sperm concentration, motility, viability and abnormality of controlled males between day 7, day 21 and day 35.

The values given are mean \pm SEM. The values with different alphabetical superscripts differ significantly ($P < 0.01-0.001$) in semen characteristics between controls and heated groups and between heated groups in days 7 (H7), 21 (H21) and 35 (H35) after the treatment (ANOVA analysis).

washed 3 times in the Hepes-HTF and stained in the CZB medium containing 20 $\mu\text{g} \cdot \text{mL}^{-1}$ bisbenzimidazole (Hoechst dye) (B2883, Sigma, Australia) with 10 $\mu\text{g} \cdot \text{mL}^{-1}$ propidium iodide (PI) (P4170, Sigma, Australia) at 37 °C. After 30–40 min, the embryos were fixed in 1% paraformaldehyde in PBS for 1 min and washed 3 times in the Hepes-HTF. The embryos were then transferred to a slide in a droplet of less than 5 μL , covered with a cover glass and squashed gently with the blunt end of a pen. The area under the cover glass was filled with the Hepes-HTF and nail varnish (Supershine, North Ryde, Australia) was used to seal around the cover glass. Finally, the differential labelled nuclei were examined using an Olympus fluorescent microscope (Japan) within a range of 330 to 385 nm of UV exciting light and of 410 to 600 nm of emission in blue. The total magnification in the eyepiece and photograph was $\times 400$ and $\times 250$, respectively. PI-labelled nuclei (dead cells/TE cells) appeared red while the Hoechst B2883 labelled nuclei (viable/ICM cells) appeared blue.

2.7. Data analysis

Comparisons of differences between groups were made using analysis of variance. A significant difference was assumed to exist when the probability of it being due to chance was less than 5%.

3. RESULTS

3.1. Semen characteristics of male mice, 7, 21 and 35 days after heat stress

Heat stress significantly reduced ($P < 0.01-0.001$) sperm concentration, motility, and viability of male mice at 7, 21 and 35 days when the greatest decrease occurred at 21 days after the treatment (Tab. I). However, the percentage of abnormal spermatozoa increased significantly ($P < 0.01-0.001$) for mice at 7, 21 and 35 days with the greatest increase observed at 21 days after the treatment (Tab. I).

3.2. The effect of paternal heat stress 7 days prior to mating on the in vivo development of mouse embryos

Paternal heat stress 7 days prior to mating did not affect the proportion of embryos developing to the zygote and 2-cell stages at 14–16 h and 34–39 h after mating, but significantly decreased the percentage of 4-cell to morulae at 61–65 h after mating and hatched blastocysts at 85–90 h after mating (Tab. II). However, the number of zygotes and morulae were higher in the heat-sired group (H7) (Figs. 1B and 1E) than in the controls (C7) (Figs. 1A and 1D) when the embryos were collected at 61–65 h and 85–90 h respectively after mating (Tab. II).

Table II. The effect of paternal heat stress 7 days prior to mating on the in vivo development of embryos.

Time of embryos recovered after mating (h)	14–16		34–39		61–65		85–90	
	C7	H7	C7	H7	C7	H7	C7	H7
Groups								
No. females	4	4	4	4	4	4	7	11
No. corpora lutea	10.8 ± 1.71	10.5 ± 1.29	8.8 ± 1.26	9.3 ± 2.22	7.8 ± 1.89	8.0 ± 0.82	8.3 ± 1.60	8.3 ± 1.56
Recovered embryos	10.3 ± 1.89	10.5 ± 1.29	8.5 ± 1.29	8.3 ± 1.50	7.8 ± 1.89	7.8 ± 0.96	7.7 ± 1.70	7.5 ± 1.64
Zygotes	9.5 ± 0.58	10.5 ± 1.29	0.5 ± 1.00	0.8 ± 0.50	0.3 ± 0.5	2.5 ± 0.82**		
2-C ^a			8.0 ± 2.16	7.3 ± 1.50	0.0 ± 0.00	0.5 ± 0.58		
4-C to M					7.5 ± 2.38	4.0 ± 2.45*	0.0 ± 0.00	1.3 ± 1.42**
B							4.9 ± 2.61	6.2 ± 2.23
HB							2.9 ± 3.20	0.0 ± 0.00***
B + HB							7.7 ± 1.60	6.2 ± 2.23*
Abnormal	0.8 ± 1.50	0.0 ± 0.00	0.0 ± 0.00	0.3 ± 0.50	0.0 ± 0.00	0.8 ± 0.50	0.0 ± 0.00	0.0 ± 0.00
Unfertilised					0.0 ± 0.00	0.3 ± 0.50		

^a C: cell, M: morula, B: blastocyst, HB: hatched blastocyst, Abnormal: abnormal embryo, Unfertilised: unfertilised ovum. The values given are mean ± SEM. The values with *, ** and *** superscripts differ significantly ($P < 0.05$ – 0.001) between the heat-sired group (H7) and controls (C7) for embryo development (ANOVA analysis).




Figure 1. A, B, C represent respectively embryos collected from the oviducts of mice at 64 h after mating with a control male (A) or one 7 (B) or 21 days (C) after heating; D, E and F represent embryos collected from the uterus of mice at 89 h after mating with a control male (D) or one 7 (E) and 21 days (F) after heating $\times 100$ (Bar = 100 μm).

3.3. The effect of paternal heat stress 21 days prior to mating on the in vivo development of mouse embryos

Paternal heat stress 21 days earlier significantly reduced the proportions of embryos developing to 2-cell, 4-cell to morulae, and prevented any embryos developing to blastocysts, but increased the number of zygotes and abnormal embryos (Tab. III and Figs. 1C and 1F). Moreover, there were more 2-cell embryos at 85–90 h after mating of heated males (Tab. III).

3.4. The effect of paternal heat stress 35 days prior to mating on the in vivo development of mouse embryos

Paternal heat stress 35 days prior to mating significantly increased the number of zygotes, but reduced the proportion of embryos developing to the 2-cell stage at 34–39 h after mating (Tab. IV). Furthermore, while there was no difference in the total number of blastocysts (B + EB), the proportion of embryos developing to blastocysts (B) was higher and fewer of those developed to expanded blastocysts (EB) in H35 than in C35 (Tab. IV).

3.5. The effect of paternal heat stress 7 and 35 days prior to mating on the number of TE and ICM in the blastocysts

There were significantly fewer total cells in the embryos sired by heat-stressed males on 7 days after heating, and in the number of cells in the trophectoderm and the inner cell mass, but, no change in the ICM/TE ratio (Tab. V). However, these differences were not significant in the embryos sired 35 days after heating (Tab. V). No blastocysts were available in the H21 group.

4. DISCUSSION

The results of this study support previous reports which have indicated that both the

whole body heating and the local heating of the scrotum decreased sperm concentration, motility and viability, but increased the number of abnormal spermatozoa in the mouse [32, 33], bull [34–36], ram [12, 37] and boar [38, 39].

The results of this study also confirmed previous reports which have shown that when spermatozoa are subjected to an elevated temperature in the uterus or in vitro or when males are exposed to high ambient temperatures, the in vivo developmental rates of preimplantation embryos are reduced in the rabbit [10, 11] and in the mouse [7, 40]. The present finding that the number of 4-cell embryos to morulae 61–65 h after mating was significantly lower in the experimental groups at 7 and 21 days after heat stress, with a higher proportion of the embryos at the zygote or 2-cell stages was different from that of previous studies which suggested that paternal heat stress significantly increases the number of 4-cell embryos [7]. This discrepancy could be attributed to the different mouse strains used, heat-stress conditions and the time of collecting embryos. However, the decrease in the number of blastocysts 85–90 h after mating is in accord with earlier results [40]. Furthermore, a common feature in this experiment and previous studies [7, 18, 41] is that the heat-stressed male can reduce the developmental rate of preimplantation embryos and increase the number of retarded embryos.

Spermatozoa ejaculated on 7 days after heating would have been in the epididymis at the time of heating [41, 42]. Although the epididymis has previously been reported to be resistant to heat [43], heat stress could disturb epididymal maturation of the spermatozoa because heat stress has been shown to damage the chromatin structure of sperm in the epididymis of the mouse [44] and the bull [45]. The results of this study have shown that heat stress significantly reduced sperm concentration, motility, viability, but increased the number of abnormal spermatozoa. A reduction in sperm quality resulted in a decrease of the developmental rate of

Table III. The effect of paternal heat stress 21 days prior to mating on the in vivo development of embryos.

Time of embryos recovered after mating (h)	14–16		34–39		61–65		85–90	
	C21	H21	C21	H21	C21	H21	C21	H21
Groups								
No. females	4	5	5	6	5	5	6	7
No. corpora lutea	10.8 ± 1.71	9.8 ± 1.30	9.2 ± 1.48	8.7 ± 1.21	8.0 ± 1.22	8.6 ± 1.82	7.7 ± 1.37	9.0 ± 2.16
Recovered embryos	10.8 ± 1.71	9.2 ± 1.79	9.0 ± 1.22	8.3 ± 1.21	7.8 ± 1.48	8.2 ± 2.59	7.3 ± 1.03	8.3 ± 1.98
Zygotes	10.8 ± 1.71	8.8 ± 1.79	0.2 ± 0.45	3.3 ± 3.20***	0.0 ± 0.00	4.2 ± 3.35***	0.0 ± 0.00	1.6 ± 0.77***
2-C ^a			8.4 ± 1.95	4.5 ± 3.21***	0.0 ± 0.00	0.6 ± 0.89	0.0 ± 0.00	1.1 ± 0.90**
4-C to M					7.8 ± 1.48	0.2 ± 0.45***	2.0 ± 2.53	0.1 ± 0.38***
B							5.7 ± 2.73	0.0 ± 0.00***
Abnormal			0.6 ± 0.89	0.5 ± 0.84	0.0 ± 0.00	3.2 ± 1.64***	0.0 ± 0.00	5.4 ± 2.36***
Unfertilised	0.0 ± 0.00	0.4 ± 0.55						

^a C, M, B, abnormal and unfertilised are the same as in Table II.

The values given are mean ± SEM. The values with ** and *** superscripts differ significantly ($P < 0.01$ – 0.001) between the heat-sired group (H21) and controls (C21) for embryo development (ANOVA analysis).

Table IV. The effect of paternal heat stress 35 days prior to mating on the in vivo development of embryos.

Time of embryos recovered after mating (h)	14–16		34–39		61–65		85–90	
	C35	H35	C35	H35	C35	H35	C35	H35
Groups								
No. females	4	4	4	5	4	4	6	5
No. corpora								
lutea	11.0 ± 2.16	10.8 ± 0.96	8.3 ± 0.96	8.8 ± 2.17	7.5 ± 1.29	7.3 ± 1.89	9.5 ± 1.38	8.6 ± 1.14
Recovered								
embryos	11.0 ± 2.16	10.3 ± 0.50	8.3 ± 0.96	8.4 ± 2.07	6.3 ± 1.26	6.3 ± 2.22	9.5 ± 1.38	8.4 ± 1.14
Zygotes	9.5 ± 0.58	10.3 ± 0.50	0.0 ± 0.00	2.4 ± 2.07***	0.0 ± 0.00	0.0 ± 0.00	0.0 ± 0.00	0.0 ± 0.00
2-C ^a			8.0 ± 0.82	5.6 ± 4.28***				
4-C to M					5.8 ± 0.96	6.3 ± 2.22	1.0 ± 1.55	0.6 ± 0.89
B							3.5 ± 3.21	5.6 ± 4.04**
EB							5.0 ± 4.73	2.2 ± 2.20**
B+EB							8.5 ± 2.88	7.8 ± 1.64
Abnormal	1.5 ± 1.73	0.0 ± 0.00	0.3 ± 0.50	0.4 ± 0.89	0.3 ± 0.50	0.0 ± 0.00	0.0 ± 0.00	0.0 ± 0.00

^a C, M, B and abnormal as defined in Table II; EB: expanded blastocysts.

The values given are mean ± SEM. The values with ** and *** superscripts differ significantly ($P < 0.01$ – 0.001) between the heat-sired group (H35) and controls (C35) for embryo development (ANOVA analysis).

Table V. The number of cells in blastocysts sired either 7 or 35 days after heating.

Sire	No. B ^a	ICM	TE	Total No. TE + ICM	Ratio (ICM/TE)
Control (C7)	20	18.2 ± 2.92	25.2 ± 6.17	43.8 ± 7.08	0.75 ± 0.19
Heated (H7)	20	14.8 ± 5.30*	19.0 ± 6.27**	33.7 ± 10.58***	0.82 ± 0.33
Control (C35)	21	17.1 ± 4.81	26.3 ± 8.35	43.5 ± 12.65	0.66 ± 0.12
Heated (H35)	17	16.4 ± 4.37	22.9 ± 5.37	39.4 ± 8.68	0.73 ± 0.17

^a B: blastocysts, ICM: inner cell mass, TE: trophoctoderm.

The values given in ratio are the sum of the addition of ICM/TE ratio in each embryo divided by the total number of blastocysts used. Values with *, ** and *** superscripts differ significantly ($P < 0.02$, 0.005 and 0.001) between the heat-sired group (H7) and controls (C7) for the number of ICM and TE cells (ANOVA analysis).

preimplantation embryos arisen from the epididymal sperm of heat-stressed males. This is in agreement with an increased mortality of embryos produced with immature, epididymal spermatozoa collected from rams during scrotal insulation [12].

Paternal heat stress 7 days prior to mating reduced the number of TE and ICM cells in the blastocyst although the ratio of ICM/TE was not different. A reduction of TE and ICM in the blastocyst could decrease its autocrine capacity which could be extremely important in any communication between the conceptus and the uterus. This could directly affect the subsequent implantation and pregnancy and it is relevant that paternal heat stress has been reported to reduce implantation rates [7, 40] and fetus weights and litter sizes [34, 46] in the mouse, and pregnancy rates in sheep [12], rats [18], pigs [6] and humans [24].

Twenty-one days after heating, there are significant reductions in the developmental rate of 2-cell embryos and an increase in the number of abnormal embryos. Additionally, no embryos developed into blastocysts when sired 21 days after heating. The increase of abnormal embryos in this study was similar to previous results published for the rat at 15 days after heating [18], and with the reduced implantation rate reported in the mouse at 21 days after heating [8]. A model proposed by Haig and Westoby [47] pre-

dicts that paternal imprinted genes will promote growth and maternal genes will inhibit growth. A paternal imprinted gene encoding U2afp-rs, an RNA processing protein is transiently transcribed at the 2-cell stage in the mouse [48]. One possibility is that heat stress could impair and/or disturb the paternal imprinted genes during spermatogenesis, thus inhibiting subsequent embryo development. Furthermore, the higher proportion of 2-cell embryos 85–90 h after mating could be that endogenous markers of zygotic gene activation including a transcription requiring complex (TRC) [49–51], HSP70.1 [52–54] and a translation initiation factor (eIF-4C) [55] might either not be transcribed transiently and/or be abnormally transcribed during the 2-cell stage. These findings merit further investigation.

Spermatozoa ejaculated 21 days after heating would have been pachytene and diplotene spermatocytes at the time of heating [42, 43] and these cells have been previously reported to be seriously damaged by heat exposure [45, 56, 57] and this study. Apparently this damage can extend to an effect on the development of preimplantation embryos.

Paternal heat stress 21 and 35 days prior to mating also increased the number of zygotes at 34–39 h, 61–65 h and 85–90 h after mating, compared with the control groups. This may be because paternal heat stress could

damage nuclear remodelling and pronuclear formation [58, 59], inhibit the first round of DNA replication [60] and/or impair the regulatory balance between cell death genes and cell survival genes [61]. Furthermore, paternal heat stress 35 days earlier reduced the number of 2-cell embryos at 34–39 h after mating and expanded blastocysts 85–90 h after mating. Since sperm ejaculated 35 days after heating would have been spermatogonia at the time of heating, spermatogonia in the testis are also damaged by heat insult because heat stress reduced sperm concentration and motility and increased the number of abnormal spermatozoa of male mice. This was in agreement with a decreased sperm concentration and the increased number of abnormal spermatozoa recovered from the cauda epididymis of male mice, 35 days after exposure to a 40 °C water bath for 2 h [32].

In conclusion, the present results further demonstrated that embryo quality is related to sperm quality. These results suggest that heat stress damaged germ cells from both the epididymis and the testis, and significantly damaged spermatocytes, and possibly also spermatogonia because there were low sperm concentration, motility, viability and high abnormality in male mice after the treatment. A reduction in sperm quality after paternal heat stress leads to a decrease in the development rate of preimplantation embryos, significantly reducing the number of 2-cell embryos, 4-cell to morula and blastocysts.

ACKNOWLEDGEMENTS

We are grateful for the assistance of Mr Jakrit Yaeram, Mr Hamish Hamilton, Mrs Lindy Mcphee, Ms Natasha Penno, Dr Sarah Robertson and Dr Graham Webb. This work was supported by the Australian Research Council, and by the award of the University of Adelaide, an Overseas Postgraduate Research Scholarship to Bi-ke Zhu.

REFERENCES

- [1] Kitayama T. Study on testicular temperature in man. *Acta Urol Japon* 1965, 11: 435–437.
- [2] Waites GMH. Temperature regulation and the testis. In: John AD, Gomes WR, VanDemark NL (Eds), *The Testis I*, Academic Press, London, 1970, p 241–279.
- [3] Waites GMH. Temperature regulation and fertility in male and female mammals. *Isr J Med Sci* 1976, 9: 982–993.
- [4] Setchell BP. The scrotum and thermoregulation. In: Setchell BP (Ed), *The Mammalian Testis*, Paul Elek Press, London, 1978, p 90–108.
- [5] Moule GR, Waites GMH. Seminal degeneration in the ram and its relation to the temperature of the scrotum. *J Reprod Fertil* 1963, 5: 433–446.
- [6] Wettemann RP, Bazer FW. Influence of environmental temperature on prolificacy of pigs. *J Reprod Fertil* 1985, 33 (Suppl): 199–208.
- [7] Bellve AR. Viability and survival of mouse embryos following parental exposure to high temperature. *J Reprod Fertil* 1972, 30: 71–81.
- [8] Burfening PJ, Elliot DS, Eisen EJ, Ulberg LC. Survival of embryos resulting from spermatozoa produced by mice exposed to elevated ambient temperature. *J Anim Sci* 1970, 30: 578–582.
- [9] Zhu B, Walker SK, Setchell BP, Maddocks S. The effect of paternal heat stress on development in vitro of preimplantation embryos in the mouse. *Proc Aust Soc Reprod Biol* 1999, 30: 62.
- [10] Howarth B, Alliston CW, Ulberg LC. Importance of uterine environment on rabbit sperm prior to fertilization. *J Anim Sci* 1965, 24: 1027–1032.
- [11] Burfening PJ, Ulberg LC. Embryonic survival subsequent to culture of rabbit spermatozoa at 38 °C and 40 °C. *J Reprod Fertil* 1968, 15: 87–92.
- [12] Mieusset R, Quintana Casares P, Sanchez Partida G, Sowerbutts SF, Zupp JL, Setchell BP. Effects of heating the testes and epididymides of ram by scrotal insulation on fertility and embryonic mortality in ewes inseminated with frozen semen. *J Reprod Fertil* 1992, 94: 337–343.
- [13] Dutt RH, Simpson EC. Environmental temperature and fertility of Southdown rams early

- in the breeding season. *J Anim Sci* 1957, 16: 136–143.
- [14] Fowler DG, Dun RB. Skin folds and merino breeding. 4. The susceptibility of rams selected for a high degree of skin wrinkle to heat induced infertility. *Aust J Exp Agric Anim Husb* 1966, 6: 121–127.
- [15] Howarth B. Fertility in the ram following exposure to elevated ambient temperature and humidity. *J Reprod Fertil* 1969, 19: 179–183.
- [16] Rathore AK. Fertility of rams heated for 1, 2, 3, and 4 days, mated to superovulated ewes. *Aust J Agric Res* 1970, 21: 355–358.
- [17] Braden AWH, Mattner PE. The effects of scrotal heating in the ram on semen characteristics, fecundity and embryonic mortality. *Aust J Agric Res* 1970, 12: 509–518.
- [18] Setchell BP, D'Occhio MJ, Hall MJ, Laurie MS, Tucker MJ, Zupp JL. Is embryonic mortality increased in normal females mated to subfertile males? *J Reprod Fertil* 1988, 82: 567–574.
- [19] Levine RJ, Bordson BL, Mathew RM, Brown MH, Stanley JM, Starr TB. Deterioration of semen quality during summer in New Orleans. *Fertil Steril* 1988, 49: 900–907.
- [20] Levine RJ, Mathew RW, Brandon Chenault C, Brown MH, Hurtt ME, Bentley KS, Mohr KL, Working PK. Differences in the quality of semen in outdoor workers during summer and winter. *N Engl J Med* 1990, 323: 12–16.
- [21] Tjoa WS, Smolensky MH, Hsi BP, Steinberger E, Smith KD. Circannate rhythm in human sperm count revealed by serially independent sampling. *Fertil Steril* 1982, 38: 454–459.
- [22] Gyllenborg J, Skakkebaek NE, Nielsen NC, Keiding N, Giwercman A. Secular and seasonal changes in semen quality among young Danish men: a statistical analysis of semen samples from 1927 donor candidates during 1977–1995. *Int J Androl* 1999, 22: 28–36.
- [23] Robinson D, Rock J. Intrascrotal hyperthermia induced by scrotal insulation: effect of spermatogenesis. *Obstet Gynecol* 1967, 29: 217–223.
- [24] Thonneau P, Ducot B, Bujan L, Mieusset R, Spira A. Effect of male occupational heat exposure on time to pregnancy. *Int J Androl* 1997, 20: 274–278.
- [25] Bamba K. Evaluation of acrosomal integrity of boar spermatozoa by bright-field microscopy using an eosin-nigrosin stain. *Theriogenology* 1988, 29: 1245–1251.
- [26] Gangrade BK, Dominic CL. Oestrous induction in unisexually grouped mice by multiple short-term exposures to males. *Biol Reprod* 1984, 31: 89–96.
- [27] Jemiolo B, Harvey S, Novotny M. Promotion of the Whitten effect in female mice by synthetic analogs of male urinary constituents. *Proc Natl Acad Sci USA* 1986, 83: 4576–4579.
- [28] Zhu B, Walker SK, Maddocks S. Optimisation of in vitro culture conditions in B6CBF1 mouse embryos. *Reprod Nutr Dev* 2004, 44: 219–231.
- [29] Quinn P, Kerin JF, Wanes GM. Improved pregnancy rate in human in vitro fertilization with the use of a medium based on the composition of human tubal fluid. *Fertil Steril* 1985, 44: 493–498.
- [30] Nicolson GL, Yanagamachi R, Yanagamachi H. Ultrastructural localisation of lectin-binding sites on the zonae pellucidae and plasma membranes of mammalian eggs. *J Cell Biol* 1975, 66: 263–274.
- [31] Chatot CL, Ziomek CA, Bavister BD, Lewis JL, Torres I. An improved culture medium supports development of random-bred 1-cell mouse embryos in vitro. *J Reprod Fertil* 1989, 86: 679–688.
- [32] Cairnie AB, Leach KE. Quantitative studies of cytological damage in mouse testis produced by exposure to heat. *Can J Genet Cytol* 1980, 22: 93–102.
- [33] Jannes P, Spiessens C, Van der Auwera I, D'Hooghe T, Verhoeven G, Vanderschueren D. Male subfertility induced by acute scrotal heating affects embryo quality in normal female mice. *Hum Reprod* 1998, 13: 372–375.
- [34] Casady RB, Myers RM, Legates JE. The effect of exposure to high ambient temperature on spermatogenesis in the dairy bull. *J Dairy Sci* 1953, 36: 14–23.
- [35] Skinner JD, Louw GN. Heat stress and spermatogenesis in *Bos indicus* and *Bos Taurus* cattle. *J Appl Physiol* 1966, 21: 1784–1790.
- [36] Vogler CJ, Bame JH, DeJarnette JM, McGilliard ML, Saacke RG. Effects of elevated testicular temperature on morphology characteristics of ejaculated spermatozoa in the bovine. *Theriogenology* 1993, 40: 1207–1219.
- [37] Smith JF. The effect of temperature on characteristics of semen in rams. *Aust J Agric Res* 1971, 22: 481–490.

- [38] Larsson K, Einarsson S. Seminal changes in boars after heat stress. *Acta Vet Scand* 1984, 25: 57–66.
- [39] Malmgren L, Larsson K. Semen quality and fertility after heat stress in boars. *Acta Vet Scand* 1984, 25: 425–435.
- [40] Bellve AR. Development of mouse embryos with abnormalities induced by parental heat stress. *J Reprod Fertil* 1973, 35: 393–403.
- [41] Oakberg EF. Duration of spermatogenesis in the mouse. *Nature* 1957, 180: 1137–1138.
- [42] Dadoune JP, Alfonsi MF. Autoradiographic investigation of sperm transit through the male mouse genital tract after tritiated thymidine incorporation. *Reprod Nutr Dev* 1984, 24: 927–935.
- [43] Vogler CJ, Saacke RG, Bame JH, Dejarnette JM, McGilliard ML. Effects of scrotal insulation on viability characteristics of cryopreserved bovine semen. *J Dairy Sci* 1991, 74: 3827–3835.
- [44] Sailer BL, Sarhar LJ, Bjordahl JA, Jost LK, Evenson DP. Effects of heat stress on the mouse testicular cells and sperm chromatin structure. *J Androl* 1997, 18: 294–301.
- [45] Karabinus DS, Vogler CJ, Saacke RG, Evenson DP. Chromatin structural changes in sperm after scrotal insulation of Holstein bulls. *J Androl* 1997, 18: 549–555.
- [46] Setchell BP, Ekpe G, Zupp JL, Surani MAH. Transient retardation in embryo growth in normal female mice made pregnant by males whose testes had been heated. *Hum Reprod* 1998, 13: 342–347.
- [47] Haig D, Westoby M. Parent-specific gene expression and the triploid endosperm. *Am Nat* 1989, 134: 147–155.
- [48] Latham KE, Rambhatla L, Hayashizaki Y, Chapman VM. Stage-specific induction and regulation by genomic imprinting of the mouse *U2afbp-rs* gene during preimplantation development. *Dev. Biol* 1995, 168: 670–676.
- [49] Latham KE, Garrels JI, Chang C, Solter D. Quantitative analysis of protein synthesis in mouse embryos I. Extensive reprogramming at the one- and two-cell stages. *Development* 1991, 112: 921–932.
- [50] Poueymirou WT, Schultz RM. Regulation of mouse preimplantation development: inhibition of synthesis of proteins in the two-cell embryo that require transcription by inhibitors of cAMP-dependent protein kinase. *Dev Biol* 1989, 133: 588–599.
- [51] Schultz RM. Regulation of zygotic gene activation in the mouse. *BioEssays* 1993, 15: 531–538.
- [52] Bensuade O, Babinet C, Morange M, Jacob F. Heat shock proteins, first major products of zygotic gene activity in mouse embryo. *Nature* 1983, 305: 331–332.
- [53] Christians E, Michel E, Adenot P, Mezger V, Rallu M, Morange M, Renard JP. Evidence for the involvement of mouse heat shock factor 1 in the atypical expression of the HSP70.1 heat shock gene during mouse zygotic genome activation. *Mol Cell Biol* 1997, 17: 778–788.
- [54] Thompson EM, Legouy EL, Christians E, Renard JP. Progressive maturation of chromatin structure regulates HSP70.1 gene expression in the preimplantation mouse embryo. *Development* 1995, 121: 3425–3427.
- [55] Davis WJ, DeSousa PA, Schultz RM. Transient expression of translation initiation factor eIF-4C during the 2-cell stage of the preimplantation mouse embryo: identification by mRNA differential display and the role of DNA replication in zygotic gene activation. *Dev Biol* 1996, 174: 190–201.
- [56] De Vita R, Calugi A, Forte D, Chiarantano C, Uccelli R, Vizzone A, Mauro F. Cellular effects of heat on mouse spermatogenesis. In: Burger G, Ploem JS, Goertler K (Eds), *Clinical Cytometry and Histometry*, Academic Press, London, 1987, p 300–302.
- [57] De Vita R, Calugi A, Chiarantano C, Forte D, Mauro F, Uccelli R. Effects of heat on mouse spermatogenesis monitored by flow cytometry. *Int J Hyperthermia* 1990, 6: 543–551.
- [58] Worrad DM, Schultz RM. Regulation of gene expression in the preimplantation mouse embryo: temporal and spatial patterns of expression of the transcription factor SP1. *Mol Reprod Dev* 1997, 46: 268–277.
- [59] Latham KE. Mechanisms and control of embryonic genome activation in mammalian embryos. *Int Rev Cytol* 1999, 193: 71–124.
- [60] Forlani S, Bonnerot C, Capgras S, Nicolas JF. Relief of a repressed gene expression state in the mouse 1-cell embryo requires DNA replication. *Development* 1998, 125: 3153–3166.
- [61] Jurisicova A, Latham KE, Casper RF, Varmuza S. Expression and regulation of genes associated with cell death during murine preimplantation embryo development. *Mol Reprod Dev* 1998, 51: 243–253.